

# Work and Energy


AP Physics – Mr. Hall

## WORKSHEET # 3


Name: \_\_\_\_\_

1. A spring has a spring constant of  $k = 55.0 \text{ N/m}$ . The spring is compressed a distance of 3.50 cm. What is the potential energy stored in the spring?
2. An 85 g wooden block is set up against a spring. The block rests on a smooth surface. The block is pushed into the spring, compressing it a distance of 2.0 cm and then released. The spring constant is  $k = 78 \text{ N/m}$ . What is the speed of the block when it reaches its initial position (where the spring was not compressed)?
3. A roller coaster starts at some height that you do not know. It goes down this hill and then goes up a second hill that is 28.5 m above the first drop at a speed of 22.5 m/s. So how high was the initial hill?
4. A 5.0 kg crate slides down a smooth ramp that is elevated at an angle of  $38^\circ$ . The length of the ramp is 2.0 m. What will be the speed of the crate at the bottom of the ramp?
5. A fireman runs up a 7.5 m ladder. The fireman has a mass of 52 kg and is carrying 15 kg of firefightin' gear. If the fireman developed 685 watts, how much time did it take to reach the top of the ladder?

6. A 3.00 kg mass,  $m_1$ , slides up a ramp. The angle for the ramp is  $28.0^\circ$ . The 3.00 kg mass is connected to a second mass,  $m_2$ , of 3.25 kg as shown by a light string with a frictionless pulley, &c. Coefficient of kinetic friction is 0.285. Find (a) the acceleration of  $m_1$ , (b) the kinetic energy of  $m_1$  after it has traveled 25.0 cm up the ramp, and (c) the work done on  $m_1$  to move it the 25.0 cm.



7. A 3.12 kg iron ball is suspended from the ceiling of a room by two cords as shown in the drawing. The ceiling is 2.20 m above the deck. The angle  $\theta$  is  $61.0^\circ$ . The other cord is perfectly horizontal. Find (a) The tension in both strings. (b) The potential energy of the ball relative to the deck. (c) if the horizontal spring were to break, what would be the velocity of the ball when the other cord is vertical?



8. A 47.0 kg projectile is launched with an initial speed of 72.0 m/s and an angle of  $39.8^\circ$  above the horizontal. The projectile lands on a hillside 7.15 s later. Neglect air friction. (a) What is the projectile's kinetic energy at the highest point of its trajectory? (b) What is the height of the impact point? (c) What is its total energy just before it hits the hillside?